REFERENCES

Ashley DL, O'Connor RJ, Bernert JT, Watson CH, Polzin GM, Jain RB, et al., (2010). Effect of differing levels of tobacco-specific nitrosamines in cigarette smoke on the levels of biomarkers in smokers. Cancer epidemiology, biomarkers and prevention. 19(6):1389-98.

Baker TB, Piper ME, McCarthy DE, Bolt DM, Smith SS, Kim SY et al.,. (2007). Time to first cigarette in the morning as an index of ability to quit smoking: Implications for nicotine dependence. Nicotine and Tobacco Research, 9, S555–S570.

Bansal H, Sandhu VS, Bhandari R, Sharma D. (2012). Evaluation of micronuclei in tobacco users: A study in Punjabi population. Contemporary Clinical Dentistry, 3(2), 184-187.

Benowitz NL, Jacob III, Denaro C, Jenkins R. (1991). Stable isotope studies of nicotine kinetics and bioavailability. Clin Pharmacol Ther, 49, 270–277.

Benowitz NL, Jacob P III, Ahijevych K, Jarvik M, Hall S, LeHouezec J, et al.,. (2002a). Biochemical verification of tobacco use and cessation. Nicotin.e and Tobacco Research 4, 149–159.

Benowitz NL & Jacob P. (1994). Metabolism of nicotine to cotinine studied by a dual stable isotope method. Clinical Pharmacology and Therapeutics 56, 483–493.

Blackford AL, Yang G, Hernandez-Avila M, Przewozniak K, Zatonski W, Figueiredo V, et al.,. (2006). Cotinine concentration in smokers from different countries: relationship with amount smoked and cigarette type. Cancer Epidemiol Biomarkers Prev, 15 (10), 1799-1804.

Bloching, M., Hofmann, A., Lautenschläger, C., Berghaus, A., & Grummt, T. (2000). Exfoliative cytology of normal buccal mucosa to predict the relative risk of cancer in the upper aerodigestive tract using the MN-assay. Oral Oncology, 36, 550–555.

Bohrer PL, Filho MS, Paiva RL, da Silva IL, Rados PV. (2005). Assessment of micronucleus frequency in normal oral mucosa of patients exposed to carcinogens. Acta Cytologica, 49, 265–272.

Brundtland, GH .(2000). Achieving worldwide tobacco control. JAMA, 284, 750–751.

Cairns, J. (1975). Mutation selection and the natural history of cancer. Nature 255, 197–200.

Caplash S, Chaudhary M and Kaur S. (2013). Micronucleus investigation in exfoliated buccal cells among tobacco chewers/ smokers and controls. International journal of Biology, Pharmacy and Allied Sciences, 2(1), 72-79.

Casartelli G, Bonatti S, De Ferrari M, Scala M, Mereu P, Marquarino G et al. (2000). "Micronucleus frequencies in exfoliated buccal cells in normal mucosa, precancerous lesions and squamous cell carcinoma," Analytical & Quantitative Cytology & Histology, 22, (6), 486–492.

Chandirasekar R, Suresh K, Sasikala K, Kumar BL, Venkatesan R, Ganesh GK and Jacob R. (2012). Genotoxicity assessment in smokeless tobacco users: a case–control study. Toxicology and Industrial Health, 29(2), 216–223.

Chandirasekar R, Kumar BL, Sasikala K, Jayakumar R, Suresh K, Venkatesan R, et al., (2014). Assessment of genotoxic and molecular mechanisms of cancer risk in smoking and smokeless tobacco users. Mutation Research, 767, 21–27.

Chatterjee S, Dhar S, Sengupta B et al. (2009). Cytogenetic monitoring in human oral cancers and other oral pathology: the micronucleus test in exfoliated buccal cells. Toxicology Mechanisms and Methods, 19 (6-7), 427–433.

Dhavan P, Bassi S, Stigler MH, Arora M, Gupta VK, Perry CL, Ramakrishnan L, Reddy KS. (2011). Using salivary cotinine to validate self-reports of tobacco use by Indian youth living in low - income neighborhoods. Asian Pacific Journal of Cancer Prevention, 12 (10), 2551-2554.

Etter JF, Vu Duc T, Perneger TV. (2000). Saliva Cotinine Levels in Smokers and Nonsmokers. American Journal of Epidemiology, 151, 251-258.

Etter J F, Perneger TV. (2001). Measurement of self reported active exposure to cigarette smoke. J Epidemiol Community Health, 55, 674–680.

Fagerström, K. (2012). Determinants of tobacco use and renaming the FTND to the Fagerström test for cigarette dependence. Nicotine and Tobacco Research, 14(1), 75–78.

Fendrich M, Mackesy-Amiti ME, Johnson TP, Hubbell A & Wilslar JS. (2005). Tobacco-reporting validity in an epidemiological drug use survey. Addictive Behaviors 30, 175–181.

Ferketich A K, Wee A G, Shultz J, and Wewers M E. (2007). Smokeless Tobacco Use and Salivary Cotinine Concentration. Addictive Behaviors, 32(12), 2953–2962.

Fiore MC, Bailey WC, Cohen SJ, Dorfman SF, Goldstein MG, Gritz ER, et al., (2000). Treating Tobacco use and Dependence. Clinical Practice Guideline. US Public Health Service, Rockville, MD.

Fu M, Fernandez E, Martínez-Sánchez JM, Pascual JA, Schiaffino A, Agudo A, et al.,. (2009) Salivary cotinine concentrations in daily smokers in Barcelona, Spain: a cross-sectional study. BMC Public Health, 9, 320.

Fu M, Martínez-Sánchez JM, Agudo A, Pascual JA, Ariza C, Moncada A, et al., (2012). Nicotine dependence and salivary cotinine concentration in daily smokers. European Journal of Cancer Prevention 21:96–102.

Halder A, Chakraborty T, Mandal K et al., (2004). Comparative study of exfoliated oral mucosal cell micronuclei frequency in normal, precancerous and malignant epithelium. International Journal of Human Genetics, 4(4), 257–260.

Hatsukami DK, Gust SW, Keenan RM. (1987). Physiologic and subjective changes from smokeless tobacco withdrawal. Clin Pharmacol Ther, 41, 103–7.

Heatherton TF, Kozlowski LT, Frecker RC, Fagerstrom KO. (1991). The Fagerstrom test for nicotine dependence: a review of the Fagerstrom Tolerance Questionnaire. Br J Addict, 86, 119–27.

Henningfield JE, Cohen C & Pickworth WB. (1993). Psychopharmacology of nicotine. In Nicotine Addiction: Principles and Management (Orleans CT & Slade J eds). Oxford University Press, Oxford, 24–45.

Holland, N. et al. (2008). The micronucleus assay in human buccal cells as a tool for biomonitoring DNA damage: the HUMN project perspective on current status and knowledge gaps. Mutat. Res., 659, 93–108.

Huang C L, Lin H H and Yang Y H. (2007). Smoking characteristics and saliva cotinine levels in Taiwanese smokers: gender differences. Journal of Clinical Nursing, 17; 2367.

Jaakkola MS, Ma J, Yang G, Chin MF, Benowitz NL, Ceraso M, Samet JM. (2003). Determinants of salivary cotinine concentrations in Chinese male smokers. Preventive Medicine, 36, 282-290.

Jadhav K, Gupta N, Ahmed MBR. (2011). Micronuclei: An essential biomarker in oral exfoliated cells for grading of oral squamous cell carcinoma. J Cytol, 28(1), 7–12.

Kamath VV, Anigol P, Setlur K. (2014). Micronuclei as prognostic indicators in oral cytological smears: A comparison between smokers and non-smokers. Clinical Cancer Investigation Journal, 3(1), 49-54.

Kandel DB, Schaffran C, Griesler PC, Hu MC, Davies M and Benowitz N. (2006). Salivary cotinine concentration versus self-reported cigarette smoking: Three patterns of inconsistency in adolescence. Nicotine Tob Res, 8(4), 525–537.

Karp I, O'Loughlin J, Hanley J, Tyndale RF, Paradis G. (2006) Risk factors for tobacco dependence in adolescent smokers. Tobacco control, 15(3), 199-204.

Khanna S, Purwar A, Singh NN, Sreedhar G, Singh S, Bhalla S. (2014). Cytogenetic biomonitoring of premalignant and malignant oral lesions by micronuclei assessment: A screening evaluation. Eur J Gen Dent, 3, 46-52.

Kozlowski LT, Porter CQ, Orleans CT, Pope MA, Heatherton T. (1994). Predicting smoking cessation with self-reported measures of nicotine dependence: FTQ, FTND, and HSI. Drug & Alcohol Dependence, 34, 211–6.

Lim KH, Feisul Idzwan M, Sumarni MG, Kee CC, Amal NM, Lim KK, Gurpreet K. (2012). Heaviness of Smoking Index, Number of Cigarettes Smoked and the Fagerström Test for Nicotine Dependence Among Adult Male Malaysians. Asian Pacific J Cancer Prev, 13, 343-346.

Manimunda SP, Benegal V, Sugunan AP, Jeemon P, Balakrishna N, Thennarasu K, et al., (2012). Tobacco use and nicotine dependency in a cross-sectional representative sample of 18,018 individuals in Andaman and Nicobar Islands, India. BMC Public Health, 12(1), 515.

Maziak W, Ward K D, Eissenberg T. (2006). Measuring exposure to environmental tobacco smoke (ETS): A developing country's perspective. Preventive Medicine, 42, 409–414.

McNeill AD, Jarvis MJ, West R, Russell MAH, Bryant A. (1987). Saliva cotinine as an indicator of cigarette smoking in adolescents. British Joumai of Addiction, 82, 1355-1360.

Mohanta A, Mohanty PK, Parida G. (2013). Genotoxicity of tobacco and alcohol on oral mucosal cells. European Journal of Experimental Biology, 3(2), 503-514.

Motgi AA, Chavan MS, Diwan NN, Chowdhery A, Channe PP, Shete MV. (2014). Assessment of cytogenic damage in the form of micronuclei in oral epithelial cells in patients using smokeless and smoked form of tobacco and non-tobacco users and its relevance for oral cancer. Journal of Cancer Research and Therapeutics, 10 (1), 165-170.

MR P, Guruprasad Y, Jose M, Saxena K, K D, Prabhu V. (2014). Comparative study of genotoxicity in different tobacco related habits using micronucleus assay in exfoliated buccal epithelial cells. Journal of Clinical and Diagnostic Research, 8(5), ZC21-ZC24.

Murray RP, Connett JE, Lauger GG & Voelker HT. (1993). Error in smoking measures: effects of intervention on relations of cotinine and carbon monoxide to self-reported smoking. American Journal of Public Health, 83, 1251–1257.

Naderi NJ, Darhadi S, Sarshar S. (2012). Micronucleus assay of buccal mucosa cells in smokers with the history of smoking less and more than 10 years. Indian Journal of Pathology & Microbiology, 55(4), 433-8.

Nesic J, Rusted J, Duka T, Jackson A. (2011). Degree of dependence influences the effect of smoking on cognitive flexibility Pharmacology, Biochemistry and Behaviors, 98(3):3 76-84.

Nuca C, Amariei C, Badea V, Zaharia A, Arendt C. (2011). Salivary Cotinine, Self-Reported Smoking Status and Heaviness of Smoking Index in Adults from Constanta, Romania. OHDM,10 (1), 22-31.

Nuca C, Amariei C, Badea V, Zaharia A, Bucur L, Arendt C. (2012). Salivary cotinine - biomarker of tobacco consumption in the assessment of passive smoking prevalence. FARMACIA, 60, 662-674.

Palve DH, Tupkari JV. (2008). Clinico-pathological correlation of micronuclei in oral squamous cell carcinoma by exfoliative cytology. J Oral Maxillofac Pathol, 12, 2-7.

Park S, Lee JY, Song TM, Cho SI. (2012). Age-associated changes in nicotine dependence. Public Health, 126, 482-489.

Rosin, M.P. (1992). The use of the micronucleus test on exfoliated cells to identify anti-clastogenic action in humans: a biological marker for the efficacy of chemopreventive agents. Mutat. Res. 267, 265–276.

Saeed H, Younis WH. (2012). A cytopathological study of the effect of smoking on the oral epithelial cells in relation to oral health status by the micronucleus assay. J Bagh Coll Dentistry, 24(3), 67-70.

Saran R, Tiwari RK, Reddy PP et al., (2008). Risk assessment of oral cancer in patients with pre-cancerous states of the oral cavity using micronucleus test and challenge assay," Oral Oncology, 44(4) 354–360.

Sarto F, Finotto S, Giacomelli L, Mazzotti D, Tomanin R, Levis AG. (1987). The micronucleus assay in exfoliated cells of the human buccal mucosa. Mutagenesis, 2(1), 11-7.

STEPS (2013). Non Communicable Diseases Risk Factors : STEPS Survey Nepal 2013. Nepal Health Research Council.

Stich, H.F., Curtis, J.R. & Parida, B.B. (1982). Application of the micronucleus test to exfoliated cells of high cancer risk groups: tobacco chewers. Int. J. Cancer, 30, 553–559.

Stich, H.F. & Rosin, M.P. (1983). Quantitating the synergistic effect of smoking and alcohol consumption with the micronucleus test on human buccal mucosa cells. Int. J. Cancer, 31, 305–308.

Stich, H.F. et al. (1988). Remission of oral leukoplakias and micronuclei in tobacco/betel quid chewers treated with beta-carotene and with beta-carotene plus vitamin A. Int. J. Cancer, 42, 195–199.

Subedi BK. (2007). 'Of what diseases are Nepalese people dying?' Kathmandu University Medical Journal, 1(17), 121–123.

Swan GE, Habina K, Means B, Jobe JB, Esposito JL.(1993). Saliva cotinine and recent smoking--evidence for a nonlinear relationship. Public Health Reports, 108(6),779-83.

Tolbert PE, Shy CM & Allen JW. (1992). Micronuclei and other nuclear anomalies in buccal smears: methods development. Mutatation Research, 271 ,69–77.

van Poppel G, Verhagen H, van't Veer P, van Bladeren PJ. (1993). Markers for cytogenetic damage in smokers: associations with plasma antioxidants and glutathione S-transferase mu. Cancer Epidemiology Biomarkers and Prevention, 2(5), 441-447.

Velicer WF, Prochaska JO, Rossi JS & Snow MG. (1992). Assessing outcome in smoking cessation studies. Psychological Bulletin, 111, 23–41.

Warnakulasuriya, S., Sutherland, G., & Scully, C. (2005). Tobacco, oral cancer, and treatment of dependence. Oral Oncology, 41, 244–260.

West R. (2004). ABC of smoking cessation. Assessment of dependence and motivation to stop smoking. BMJ, 328, 338-339.

WHO (2010). Global status report on noncommunicable disease. Geneva: World Health Organization.

WHO (2002). The world health report 2002, reducing risks, promoting healthy life. Geneva: World Health Organization.

Woodward M, Tunstall-Pedoe H, Smith WC & Tavendale R. (1991). Smoking characteristics and inhalation biochemistry in the Scottish population. Journal of Clinical Epidemiology 44, 1405–1410.

Wu PA, Loh CH, Hsieh LL, Liu TY, Chen CJ, Liou SH. (2004). Clastogenic effect for cigarette smoking but not areca quid chewing as measured by micronuclei in exfoliated buccal mucosal cells. Mutation Research, 562(1-2), 27-38.